arXiv:hep-th/0311272v1 28 Nov 2003

The Dark Energy and the Fate of Universe

Mian WANG

Department of Physics, Henan Normal University

Xinxiang, Henan, China 453002

e-mail: wangm@henannu.edu.cn

Abstract

Recent observations confirm that our universe is flat and consists of a dark energy component $\Omega_{DE} \simeq 0.7$. This dark energy is responsible for the cosmic acceleration as well as determines the feature of future evolution of the universe. In this paper, we discuss the dark energy of universe in the framework of scalar-tensor cosmology. It is shown that the dark energy is the main part of the energy density of the gravitational scalar field and the future universe will expand as $a(t) \sim t^{1.3}$.

PACS number(s): 98.80.Cq, 04.50.+h

Observations of large scale structure, the Hubble diagram of type Ia supernovae and the angular power spectrum of the cosmic microwave background *et al*, all indicate that the universe is flat and is undergoing cosmic acceleration by virtue of a dark energy component, $\Omega_{DE} \simeq 0.7$, with negative pressure [1]. It is supposed that the dark energy may consist of a cosmological constant (vacuum energy) or the quintessence, such as a scalar field with negative pressure [2]. It is important to investigate the nature of this energy component, because it is not only responsible for the present accelerated expansion but also will determinate the fate of our universe. For example, if it is a cosmological constant, the universe will expand exponentially in the future. On the other hand, if the dark energy is the energy of a slowly changing scalar field φ with $w_\varphi \simeq -1$ and the potential $V(\varphi)$ driving the present stage of acceleration decreases slowly and vanishes eventually, the speed of expansion decreases after a transient de Sitter-like stage and reaches Minkowski regime [2], or the potential falls to $V(\varphi) = -\infty$, the universe eventually collapse, even if it is flat[3]. In this letter we investigate the dark energy in the framework of scalar-tensor cosmology[4] with an added potential term.

The scalar-tensor cosmology We should assume a modified action of [4] by adding a $V(\phi)$ term as follows

$$\mathcal{A}=\int d^4x\sqrt{-g}\,\left[-\phi R-\frac{\omega}{\phi}\phi\cdot\phi-2\phi\lambda(\phi)-2\phi V(\phi)-\frac{\Gamma(u\cdot\phi)^2}{1-\phi}+16\pi L_m\right], \tag{1}$$

in which L_m is the Lagrangian of matter, $\phi\cdot\phi\equiv\phi_{,\sigma}\phi^{,\sigma}$, $u\cdot\phi\equiv u_\mu\phi^{,\mu}$, u_μ is the four-velocity, $\phi^{,\mu}\equiv\partial\phi/\partial x_\mu$, and Γ is a constant of mass dimension 0. The $\Gamma-$term describes the coupling of the field ϕ with matter which is described as an ideal fluid. The coupling function $\omega(\phi)$ and the cosmological function $\lambda(\phi)$ are given as

$$2\omega(\phi)+3=\frac{\xi}{1-\phi}\,, \tag{2}$$

$$\lambda(\phi)=2\xi\beta(1-\phi-\phi\;ln\,\phi), \tag{3}$$

where ξ and β are two dimensionless constants, here we set $\xi=7.5,\;\beta=1.7\times10^{-16}$ as [4] and $\Gamma=0.06$. The potential $V(\phi)$ is

$$V(\phi)=\beta^2\sqrt{1-\phi}\;\;e^{-\alpha\phi} \tag{4}$$

in which α is a constant of order $O(10^2)$, it is set to be 81.5 in the model.

Then the Friedmann equations read

$$H^2=\frac{8\pi}{3\phi}(\rho_m+\rho_\phi)-\frac{k}{a^2}\,, \tag{5}$$

$$\frac{\ddot{a}}{a}=-\frac{4\pi}{3\phi}(\rho_m+\rho_\phi+3p_m+3p_\phi)\,, \tag{6}$$

and

$$\begin{aligned}\ddot{\phi}+3H\dot{\phi}\;\;&+\;\;\frac{\dot{\phi}^2}{2(1-\phi)}+\frac{8\pi(3p_m-\rho_m)}{2\omega+3}-\frac{\Gamma}{2\omega+3}\frac{\phi\dot{\phi}^2}{(1-\phi)^2}-\frac{2\Gamma}{2\omega+3}\frac{\phi\ddot{\phi}}{1-\phi}\\ &=\;\;4\beta\phi(1-\phi^2)+\frac{2}{\xi}\phi\sqrt{1-\phi}\,\beta^2\;e^{-\alpha\phi}\left(1-\frac{1}{2}\phi+\alpha(1-\phi)\right),\end{aligned} \tag{7}$$

$$\dot{\rho}_m+3H(\rho_m+p_m)=\frac{\Gamma}{4\pi}\left(\frac{\dot{\phi}\ddot{\phi}}{1-\phi}+\frac{1}{2}\frac{\dot{\phi}^3}{(1-\phi)^2}+3H\frac{\dot{\phi}^2}{1-\phi}\right), \tag{8}$$

where ρ_m , p_m and ρ_ϕ , p_ϕ are energy density and pressure of the matter and field ϕ respectively. The expressions are

$$\rho_\phi=\frac{1}{8\pi}\left(\phi\lambda+\phi V+\frac{\omega\dot{\phi}^2}{2\phi}-3H\dot{\phi}-\frac{3}{2}\frac{\Gamma\dot{\phi}^2}{1-\phi}\right), \tag{9}$$

$$p_\phi=\frac{1}{8\pi}\left(-\phi\lambda-\phi V+\frac{\omega\dot{\phi}^2}{2\phi}+2H\dot{\phi}+\ddot{\phi}-\frac{1}{2}\frac{\Gamma\dot{\phi}^2}{1-\phi}\right). \tag{10}$$

Exponential inflation When the field ϕ rolls down the potential hill from $\phi\simeq0$, the universe expands exponentially as described in [4] since $V(\phi)$ can be neglected when $\phi<1$. The Hubble parameter H is determined by the value of $\lambda(\phi)\simeq2\xi\beta$ when $\phi\ll1$. In the duration of inflation, the energy density of the scalar field ρ_ϕ dominates and $p_\phi=-\rho_\phi$. It is evident that ρ_ϕ acts as a cosmological constant.

Power-law expansion As ϕ increases, the $\dot{\phi}^2$ term in equation(9) increases also, therefore $w_\phi \equiv p_\phi/\rho_\phi$ increases quickly. The inflation ends dynamically when $\rho + 3p \simeq \rho_\phi + 3p_\phi$ increases from $-2\rho_\phi$ to zero ($\phi \sim 1$). Then the universe begins the era of expansion according to power law $a \sim t^{2/3}$, because the energy density ρ_ϕ remains to be dominating and the scalar ϕ behaves as massive particle with effective mass $\sim 2\sqrt{\beta}$ when ϕ goes near to the bottom of potential well. That is to say, the gravitational scalar field ϕ behaves as vacuum energy in the era of inflation and as cold dark matter after inflation.

After inflation, ϕ approaches near to its bottom value 1. Let

$$\phi = 1 - \sigma^2,$$

it follows

$$(1-2\gamma)\ddot{\sigma} + 3H\dot{\sigma} + 4\beta\sigma = -sign(\sigma)\frac{\beta^2 e^{-\alpha}}{2\xi}, \tag{11}$$

in which

$$\gamma \equiv \frac{\Gamma}{\xi}.$$

From equation (11), we see that the $\sigma(t)$ oscillates with frequency ω_0:

$$\omega_0^2 = \frac{4\beta}{1-2\gamma}.$$

Then we set

$$\sigma(t) = f(t)\ \cos\omega_0 t,$$

we have equations

$$\dot{f} = -\frac{3H}{2(1-2\gamma)}f, \tag{12}$$

$$\ddot{f} = -\frac{3H}{1-2\gamma}\dot{f} - \frac{2\beta^2 e^{-\alpha}}{\pi\xi}, \tag{13}$$

and

$$\dot{\rho}_m + 3H(\rho_m + p_m) = \frac{\Gamma}{\pi}\left(\frac{3\beta^2 e^{-\alpha}}{2\pi\xi(1-2\gamma)}Hf + \frac{3\beta}{1-2\gamma}Hf^2\right). \tag{14}$$

The energy density of ϕ consists of ρ_{DM} and ρ_{DE}:

$$\rho_{DM} = \frac{\xi\beta}{2\pi}(1-2\gamma)f^2, \qquad \rho_{DE} = \frac{\beta^2 e^{-\alpha}}{4\pi^2}f \tag{15}$$

and the pressure of ϕ is

$$p_\phi = -(1-\frac{1}{\xi})\frac{2\beta^2 e^{-\alpha}}{4\pi^2}f. \tag{16}$$

We notice that in equation(15), the multiplier constants $\frac{\xi\beta}{2\pi}(1-2\gamma) = O(10^{-16})$ and $\frac{\beta^2 e^{-\alpha}}{4\pi^2} = O(10^{-69})$. Therefore, when $f > 10^{-53}$, the ρ_{DM} dominates, equation(12) has solution

$$f = \frac{1}{\sqrt{3\xi\beta}t} \tag{17}$$

and

$$H = \frac{2(1-2\gamma)}{3t}, \qquad a(t) \sim t^{\frac{2}{3}(1-2\gamma)}. \tag{18}$$

Deceleration-acceleration transition When $f(t)$ decreases to less than 10^{-53}, correspondingly $t > 10^{60}M_P^{-1}$, the dark energy dominates gradually. In this transient stage, we have not analytic solutions. From numerical solutions by computer, we have the results: the deceleration-acceleration transition occurs at $z = 0.54$. For the present universe, $t_0 = 14Gy = 8.20\times10^{60}M_P^{-1}$, $H_0 = 74km/sMpc = 1.30\times10^{-61}M_P$, it is found that

$$H(t) = \frac{1.066}{t}, \qquad a(t) \sim t^{1.066}. \tag{19}$$

The deceleration parameter $q_0 \equiv H_0^{-2}(\ddot{a}/a)_0 = -0.06191$.

The future evolution of the universe After this transition period we can find an analytic solution. In fact, when $t >> 10^{60}M_P$, the ρ_{DE} dominates in the righthand side of equation(15), then $f(t) \sim t^{-2}$, we have solutions

$$H(t) = \frac{4(1-2\gamma)}{3t} = \frac{1.312}{t}, \tag{20}$$

$$a(t) = a_0\left(\frac{t}{t_0}\right)^{1.312} \tag{21}$$

and

$$f(t) = f_0\left(\frac{t_0}{t}\right)^2, \qquad \rho_m(t) = \rho_{m0}\left(\frac{t_0}{t}\right)^2. \tag{22}$$

The deceleration parameter $q = -(n-1)/n = -0.2378$ and the equation of motion $w = -0.49$. Therefore $\Omega_m + \Omega_{DM} = (1+q)/(2-q) = 0.34$ and we have $\Omega_{DE} = 0.66$, $\Omega_{DM} = 0.3$ and $w_{DE} = -0.75$.

In conclusion, we have shown that our universe was dominated by the energy density ρ_ϕ of the gravitational scalar field with a negative pressure $p_\phi = -\rho_\phi$ in the period of inflation, afterwards then dominated by a cold dark matter $\rho_{DM} \simeq \rho_\phi$ after the inflation. After the time of $10^{60}M_p^{-1}$, the dark energy Ω_{DE} increased to accelerate the universe. Eventually our universe will expand with a power law $a(t) \sim t^{1.3}$ with $\Omega_{DE} = 0.66$ and $w_{DE} = -0.75$.

Reference

1. A.G. Riess *et al.,* Astrophys. J. **116** (1998) 1009. S. Perlmutter et al., Astrophys. J. **517** (1999) 565. C. Pryke *et al.,* Astrophys. J. **568** (2002) 46. L. Verde *et al.,* MNRAS **335** (2002) 432. N. Bahcall *et al.,* Science **284** (1999) 1481. D.N. Spergel *et al.,* Astrophys. J. Suppl. **148** (2003)175.
2. C. Wetterich, Nucl. Phys. B **302** (1988) 668. I. Zlatev *et al.,* Phys. Rev. Lett. **82** (1999) 896.
3. G.N. Felder *et al.,* Phys. Rev. D **66** (2002) 023507.
4. M. WANG, Phys. Rev. D **61** (2000) 123511.